


EXHIBIT 1

Relative Risk for Adults Age 50 and Older of Having at Least One Adverse COVID-19 Experience During the First Pandemic Wave

Odds ratios relative to US




Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All odds ratios between the US and other countries are statistically significant at the $p < .01$ level.
Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Data based on responses from 44,675 adults age 50 and older in 29 countries.

EXHIBIT 2

Adults Age 50 and Older Who Reported Being Infected or Hospitalized with COVID-19 During the First Pandemic Wave

Percent




Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All percentage differences between the US and other countries are statistically significant at the $p < .01$ level. Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Sample includes 44,675 adults age 50 and older in 29 countries.

EXHIBIT 3

Adults Age 50 and Older Who Reported Forgoing Care During the First Pandemic Wave

Percent




Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All percentage differences between the US and other countries are statistically significant at the $p < .01$ level. Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Sample includes 44,675 adults age 50 and older in 29 countries.

EXHIBIT 4

Adults Age 50 and Older Who Reported the Death of a Friend or Relative from COVID-19 During the First Pandemic Wave

Percent




Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All percentage differences between the US and other countries are statistically significant at the $p < .01$ level. Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Sample includes 44,675 adults age 50 and older in 29 countries.

EXHIBIT 5

Adults Age 50 and Older Who Reported Losing Employment During the First Pandemic Wave


Percent



Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All percentage differences between the US and other countries are statistically significant at the $p < .01$ level. Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Sample includes 44,675 adults age 50 and older in 29 countries.

EXHIBIT 6

Adults Age 50 and Older with at Least One Adverse COVID-19 Experience During the First Pandemic Wave



Notes: For this study, March through September 2020 is considered the “first pandemic wave.” All percentage differences between the US and other countries are statistically significant at the $p < .01$ level. Data: Survey of Health, Ageing, and Retirement in Europe (SHARE), Health and Retirement Study (HRS), and English Longitudinal Study of Ageing (ELSA) surveys, 2018–2019 and summer 2020. Sample includes 44,675 adults age 50 and older in 29 countries.